

### Numbers of Fish Recorded

The total fish count in 2001 at all 8 sites was 34,901, giving an average of 1,501 fish per 100 m<sup>2</sup>. At the shallower sites the density was 1,431 fish per 100 m<sup>2</sup> and at the 20m depth sites 1,628 per 100 m<sup>2</sup> (Table 4).

Although the fish were counted into six total length categories (see Methodology) the data is summarised here into three categories: 1 to 10 cm, 11 to 20 cm, and >20 cm, by transect depth. At the shallow and deep transects, the largest percentage of fish was in the 1 to 10 cm category (82% and 70%, respectively), and the smallest percentage in the 11 - 20 cm category (4% and 5%, respectively). The abundance of fishes in the 1 to 10 cm category at both transect depths was primarily due to large numbers of fish from families with numerous small-sized species (Serranidae (fairybasslets), Apogonidae, and Pomacentridae), and secondarily because of the number of juvenile life-stages (e.g. Serranidae).

Table 4. Number of fish counted, by transect depths and fish size groups, during the Aldabra Marine Programme surveys in February 2001.

	10 m Transects (Site 7 - 5m)			20 m Transects (Site 7 - 15m)			
Total Survey Area - 2325 m <sup>2</sup>	Area Surveyed: 1500 m <sup>2</sup>			Area Surveyed: 825 m <sup>2</sup>			
	Size (cm)	Size (cm)	Size (cm)	Size (cm)	Size (cm)	Size (cm)	
Family/Genus species	1 to 10	11 to 20	>20	1 to 10	11 to 20	>20	Total
<b>MURAENIDAE</b>							
<i>Gymnothorax flavimarginatus</i>			1				1
<b>BELONIDAE</b>							
<i>Tylosurus crocodilus</i>			1				1
<b>HOLOCENTRIDAE</b>							
<i>Myripristis adusta</i>		1	1		9	1	12
<i>Myripristis berndti</i>		40		17	226	14	297
<i>Myripristis murdjan</i>		22	14				36
<i>Neoniphon sammara</i>		6				1	7
<i>Sargocentron caudimaculatum</i>		31		3	58		92
<i>Sargocentron diadema</i>				13			13
<i>Sargocentron spiniferum</i>						2	2
<b>SERRANIDAE</b>							
<i>Aethaloperca rogaa</i>			2			550	552
<i>Cephalopholis argus</i>		1	11		5	4	21
<i>Cephalopholis leoparda</i>	2	1			1		4
<i>Cephalopholis miniata</i>		1	5		2	12	20
<i>Cephalopholis spiloparaea</i>	1	16	9		1		27
<i>Cephalopholis urodeta</i>	1	10			5		16
<i>Dermatolepis striolatus</i>						1	1
<i>Epinephelus fasciatus</i>	1	12	3		11		27
<i>Epinephelus fuscoguttatus</i>					1		1
<i>Epinephelus polyphekadion</i>			2			2	4
<i>Epinephelus spilotoceps</i>		1	2			1	4
<i>Gracilia albomarginata</i>		1	5			4	10
<i>Plectropomus laevis</i>						1	1

Table 4 (continued)

	10 m Transects (Site 7 - 5m)			20 m Transects (Site 7 - 15m)			
Total Survey Area - 2325 m <sup>2</sup>	Area Surveyed: 1500 m <sup>2</sup>			Area Surveyed: 825 m <sup>2</sup>			
	Size (cm)	Size (cm)	Size (cm)	Size (cm)	Size (cm)	Size (cm)	
Family/Genus species	1 to 10	11 to 20	>20	1 to 10	11 to 20	>20	Total
<i>Variola louti</i>			1			2	3
<i>Nemanthias carberryi</i>				1			1
<i>Pseudanthias cooperi</i>	3			8	6		17
<i>Pseudanthias evansi</i>	281			100			381
<i>Pseudanthias squamipinnis</i>	928	1		1098			2027
<i>Plectropomus areolatus</i>		1	2			3	6
<i>Cephalopholis nigripinnis</i>	10	37	2		15		64
<i>Pseudanthias ignitis</i>					6		6
<i>Anyperodon leucogrammicus</i>	1						1
<b>APOGONIDAE</b>							
<i>Apogon apogonoides</i>	400	2		1835			2237
<i>Cheilodipterus macrodon</i>					1		1
<b>HAEMULIDAE</b>							
<i>Plectorhinchus gaterinus</i>						7	7
<i>Plectorhinchus obscurus</i>			4			3	7
<i>Plectorhinchus orientalis</i>						4	4
<b>LUTJANIDAE</b>							
<i>Aphareus furca</i>	1		11		1	4	17
<i>Aprion virescens</i>						3	3
<i>Lutjanus bengalensis</i>				1			1
<i>Lutjanus bohar</i>	1	6	18	1	2	15	43
<i>Lutjanus gibbus</i>			6				6
<i>Lutjanus kasmira</i>					4	711	715
<i>Lutjanus monostigma</i>			5		1	8	14
<b>CAESIONIDAE</b>							
<i>Caesio xanthonota</i>			69		15	10	94
<i>Pterocaesio lativittata</i>			60		15		75
<i>Pterocaesio tile</i>			2400		1	800	3201
<b>MULLIDAE</b>							
<i>Parupeneus barbarensis</i>			1			1	2
<i>Parupeneus bifasciatus</i>	1	10	4		3	2	20
<i>Parupeneus cyclostomus</i>		3	4			3	10
<i>Parupeneus macronema</i>	148	31	3	29	7	1	219
<i>Parupeneus pleurostigma</i>		1		1		2	4
<i>Mulloidichthys vanicolensis</i>						550	550
<i>Parupeneus rubescens</i>		1					1

Table 4 (continued)

	10 m Transects (Site 7 - 5m)			20 m Transects (Site 7 - 15m)			
Total Survey Area - 2325 m <sup>2</sup>	Area Surveyed: 1500 m <sup>2</sup>			Area Surveyed: 825 m <sup>2</sup>			
	Size (cm)	Size (cm)	Size (cm)	Size (cm)	Size (cm)	Size (cm)	
Family/Genus species	1 to 10	11 to 20	>20	1 to 10	11 to 20	>20	Total
<b>CHAETODONTIDAE</b>							
<i>Chaetodon auriga</i>	4	11		9	6		30
<i>Chaetodon bennetti</i>	1			2	3		6
<i>Chaetodon falcula</i>		2		2	10		14
<i>Chaetodon guttatissimus</i>	19	3		14	3	1	40
<i>Chaetodon kleinii</i>				15	1	4	20
<i>Chaetodon lineolatus</i>						2	2
<i>Chaetodon lunula</i>		6		2	2		10
<i>Chaetodon meyeri</i>	3	1		2	1		7
<i>Chaetodon trifasciatus</i>	8	4					12
<i>Chaetodon xanthocephalus</i>	2	5	2				9
<i>Chaetodon zanzibariensis</i>					5		5
<i>Forcipiger flavissimus</i>	9	4	7	16	6		42
<i>Hemitaenichthys zoster</i>	12			7			19
<i>Heniochus acuminatus</i>		1					1
<i>Heniochus monoceros</i>					1		1
<i>Chaetodon interruptus</i>		1			1		2
<i>Chaetodon madagaskariensis</i>		3		3	2		8
<i>Chaetodon vagabundus</i>		2					2
<i>Heniochus singularis</i>			2				2
<b>LETHRINIDAE</b>							
<i>Lethrinus nebulosus</i>						3	3
<i>Monotaxis grandoculis</i>	5	21	5	8	18	2	59
<b>EPHIPPIDAE</b>							
<i>Platax orbicularis</i>			5			3	8
<b>MALACANTHIDAE</b>							
<i>Malacanthus latovittatus</i>				1			1
<b>PINGUIPEDIDAE</b>							
<i>Parapercis hexophthalma</i>	3	4		1			8
<i>Parapercis millipunctata</i>	5						5
<i>Parapercis signata</i>				6			6
<b>POMACANTHIDAE</b>							
<i>Apolemichthys trimaculatus</i>	1	7			4		12
<i>Centropyge acanthops</i>	17			28			45
<i>Centropyge multispinis</i>	59	4		37			100
<i>Pygoplites diacanthus</i>		1	2		2	2	7
<i>Centropyge bispinosa</i>	1			3			4
<i>Centropyge debelius</i>				6			6

Table 4 (continued)

	10 m Transects (Site 7 - 5m)			20 m Transects (Site 7 - 15m)			
Total Survey Area - 2325 m <sup>2</sup>	Area Surveyed: 1500 m <sup>2</sup>			Area Surveyed: 825 m <sup>2</sup>			
	Size (cm)	Size (cm)	Size (cm)	Size (cm)	Size (cm)	Size (cm)	
Family/Genus species	1 to 10	11 to 20	>20	1 to 10	11 to 20	>20	Total
<b>POMACENTRIDAE</b>							
<i>Amphiprion chrysogaster</i>					2		2
<i>Amphiprion clarkii</i>				1			1
<i>Chromis dimidiata</i>	1005			705			1710
<i>Chromis nigrura</i>	267			36			303
<i>Chromis ternatensis</i>	565			164			729
<i>Chromis weberi</i>	374	21		36			431
<i>Dascyllus aruanus</i>	2			10			12
<i>Dascyllus carneus</i>				63			63
<i>Dascyllus trimaculatus</i>	1			14			15
<i>Lepidozygus tapeinosoma</i>	11695			4290			15985
<i>Plectroglyphidodon dickii</i>	14	5		5			24
<i>Plectroglyphidodon johnstonianus</i>	49			4			53
<i>Plectroglyphidodon lacrymatus</i>	91	8		5			104
<i>Pomacentrus caeruleus</i>	58			11			69
<i>Pomacentrus sulfureus</i>	1						1
<i>Chromis xutha</i>				107			107
<i>Abudefduf sexfasciatus</i>	2						2
<i>Amphiprion allardi</i>				2	3		5
<i>Chromis atripectoralis</i>	1						1
<i>Pomacentrus chrysurus</i>	1						1
<i>Pomacentrus philippinus</i>				1			1
<b>LABRIDAE</b>							
<i>Anampses meleagrides</i>	4	2		4	6		16
<i>Bodianus axillaris</i>	5	1	1	2	5		14
<i>Bodianus bilunulatus</i>			1		1	1	3
<i>Bodianus diana</i>		5	1	3	8	3	20
<i>Cirrhilabrus exquisitus</i>	604	12		138			754
<i>Coris cuvieri</i>		2					2
<i>Coris freiei</i>		4	1			2	7
<i>Epibulus insidiator</i>	1						1
<i>Gomphosus caeruleus</i>	25	16		8	5		54
<i>Halichoeres cosmetus</i>	134	9		49	6		198
<i>Halichoeres hortulanus</i>	2	8		1	8		19
<i>Halichoeres vrolikii</i>	3	3					6
<i>Hemigymnus fasciatus</i>		1	1				2
<i>Hologymnosus doliatus</i>		2		1			3
<i>Labroides bicolor</i>	9	3		5	1		18
<i>Labroides dimidiatus</i>	104			110			214
<i>Labrobsis xanthonata</i>	1			6	1		8
<i>Macropharingodon bipartitus</i>	11	2					13
<i>Pseudocheilinus evanidus</i>	61			94			155
<i>Pseudocheilinus hexataenia</i>	174			47			221

Table 4 (continued)

	10 m Transects (Site 7 - 5m)			20 m Transects (Site 7 - 15m)			
Total Survey Area - 2325 m <sup>2</sup>	Area Surveyed: 1500 m <sup>2</sup>			Area Surveyed: 825 m <sup>2</sup>			
	Size (cm)	Size (cm)	Size (cm)	Size (cm)	Size (cm)	Size (cm)	
Family/Genus species	1 to 10	11 to 20	>20	1 to 10	11 to 20	>20	Total
<i>Pseudocheilinus octotaenia</i>	31	7		3	1		42
<i>Pseudodax molucanus</i>		2					2
<i>Thalassoma amblycephalum</i>	15	2	1	20	3		41
<i>Thalassoma hardwicke</i>		2		2			4
<i>Thalassoma herbraicum</i>	14	33	4	4	9	2	66
<i>Thalassoma lunare</i>	6	1		1	18		26
<i>Anampses caeruleopunctatus</i>			1				1
<i>Bodianus anthioides</i>			1		3		4
<i>Bodianus mesothorax</i>		1					1
<i>Coris batuensis</i>	9	9	1	16	2		37
<i>Hologymnosus annulatus</i>	1						1
<i>Novaculichthys taeniourus</i>	1	2	1		2		6
<i>Pseudojuloides kaleidas</i>				14			14
<i>Stethojulis albivittata</i>	8	1		1	1		11
<b>CIRRHITIDAE</b>							
<i>Cirrhitichthys oxycephalus</i>	10			11			21
<i>Paracirrhites arcatus</i>	28	1		10			39
<i>Paracirrhites forsteri</i>	7			10	1		18
<b>SCARIDAE</b>							
<i>Scarus sordidus</i>	10	63	23	4	14	3	117
<i>Scarus rubroviolatus</i>						1	1
<i>Scarus frenatus</i>			2			1	3
<i>Scarus strongylocephalus</i>		1	2			5	8
<b>CARANGIDAE</b>							
<i>Caranx ignobilis</i>			22			3	25
<i>Caranx melampygus</i>			32			18	50
<i>Caranx sexfasciatus</i>						515	515
<i>Elagatis bipinnulata</i>						2	2
<b>BLENNIIDAE</b>							
<i>Aspidontus taeniatus</i>	6	1		8			15
<i>Ecsenius midas</i>				4			4
<i>Plagiotremus rhinorhynchus</i>				1			1
<i>Plagiotremus tapeinosoma</i>	6			3			9
<i>Cirripectes castaneus</i>	16	1					17
<b>GOBIIDAE</b>							
<i>Valenciennea helsdingeni</i>	2						2
<i>Valenciennea strigata</i>	4						4

Table 4 (continued)

	10 m Transects (Site 7 - 5m)			20 m Transects (Site 7 - 15m)			
Total Survey Area - 2325 m <sup>2</sup>	Area Surveyed: 1500 m <sup>2</sup>			Area Surveyed: 825 m <sup>2</sup>			
	Size (cm)	Size (cm)	Size (cm)	Size (cm)	Size (cm)	Size (cm)	
Family/Genus species	1 to 10	11 to 20	>20	1 to 10	11 to 20	>20	Total
<b>ACANTHURIDAE</b>							
<i>Acanthurus auranticavus</i>			2				2
<i>Acanthurus leucocheilus</i>		6			9	6	21
<i>Acanthurus leucosternon</i>	11	76	1	1	4		93
<i>Acanthurus thompsoni</i>		5	5	11	67		88
<i>Ctenochaetus striatus</i>	75	30	7	11	3	2	128
<i>Ctenochaetus strigosus</i>	156	147		48	22		373
<i>Naso brevirostris</i>			55		2		57
<i>Naso lituratus</i>			3			1	4
<i>Zebrasoma scopas</i>	32	4		15	4		55
<i>Acanthurus lineatus</i>		4	1				5
<i>Acanthusus nigricauda</i>			4				4
<i>Acanthurus tennenti</i>		1					1
<i>Naso vlamingii</i>						14	14
<i>Zebrasoma desjardini</i>		1			1		2
<b>ZANCLIDAE</b>							
<i>Zanclus cornutus</i>	1	6			4		11
<b>BALISTIDAE</b>							
<i>Balistapus undulatus</i>		2			3		5
<i>Balistoides viridescens</i>			1				1
<i>Melichthys indicus</i>		15	48		3	6	72
<i>Odonus niger</i>		3	7				10
<i>Sufflamen bursa</i>		3			1		4
<i>Sufflamen chrysopterus</i>	5	5		3	1		14
<b>OSTRACIIDAE</b>							
<i>Ostracion meleagris</i>	1			1			2
<b>TETRAODONTIDAE</b>							
<i>Arothron meleagris</i>	1					2	3
<i>Arothron nigropunctatus</i>		2	1		2	1	6
<i>Canthigaster valentini</i>	7						7
<i>Canthigaster coronata</i>	1						1
<b>SCORPAENIDAE</b>							
<i>Scorpaenopsis diabolus</i>		1					1
<b>SIGANIDAE</b>							
<i>Siganus stellatus</i>			2				2

Table 4 (continued)

	10 m Transects (Site 7 - 5m)			20 m Transects (Site 7 - 15m)			
Total Survey Area - 2325 m <sup>2</sup>	Area Surveyed: 1500 m <sup>2</sup>			Area Surveyed: 825 m <sup>2</sup>			
	Size (cm)	Size (cm)	Size (cm)	Size (cm)	Size (cm)	Size (cm)	
Family/Genus species	1 to 10	11 to 20	>20	1 to 10	11 to 20	>20	Total
<b>MOBULIDAE</b>							
<i>Manta briostriis</i>						1	1
<b>MICRODESMIDAE</b>							
<i>Nemateleotris magnifica</i>	46			22			68
<i>Ptereleotris evides.</i>		2		3			5
<b>MONACANTHIDAE</b>							
<i>Amanses scopas</i>		2			2		4
<i>Cantherinus pardalis</i>		1					1
<b>Total Fish Counted</b>	<b>17702</b>	<b>863</b>	<b>2901</b>	<b>9418</b>	<b>689</b>	<b>3328</b>	<b>34901</b>
<b>Number of Species</b>	<b>86</b>	<b>92</b>	<b>62</b>	<b>84</b>	<b>74</b>	<b>57</b>	<b>191</b>
<b>Fish/100m<sup>2</sup></b>	<b>12</b>	<b>0.6</b>	<b>2</b>	<b>11</b>	<b>0.8</b>	<b>4</b>	<b>1501</b>

### *Fish Species Distribution*

The pattern of fish species distribution around the outer reef of the atoll in February 2001 (Figure 22), followed the same general pattern as in November 1999 (Figure 23). Sites 5, 3, 1, and 4 on the eastern portion of the atoll ranked 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> and 4<sup>th</sup>, respectively, for the least number of species in both years. Sites 2, 6, 7, and 8 (2001 only) on the western portion of the atoll accounted for the most number of species of fish on the transects in both years.

The difference in the species numbers from one end of the atoll to the other is striking. Thirty species were counted at Site 5 compared to a range of 62 to 115 species at the other sites. Furthermore, there were significant correlations between the number of species of fish surveyed on both the 10 m depth transects ( $R^2 = 0.41$ ,  $F_{0.05,1} = 4.16$ ,  $P = 0.087$ ) and the 20 m depth transects ( $R^2 = 0.47$ ,  $F_{0.05,1} = 5.23$ ,  $P = 0.062$ ) at Sites 1-8, and their relative positions from east to west along the shoreline of Aldabra Atoll (Figure 22).

### *Fish Numbers Distribution*

In February 2001, the number of fish per 100 m<sup>2</sup>, combining shallow and deep transects ranged from 299 fish (62 species) at Site 3, to 3,867 fish (89 species) at Site 6 (Figure 22). The 550 fish per 100 m<sup>2</sup> at Site 5 is unusually high for the low number of species at this site and was due to a school of 800 large fusilier, *Pterocaesio tile* (Caesionidae) crossing the transect line. The numbers of the fish counted in both shallow and deep transects combined, at Sites 1, 2, 3, 4, 6, and 7 were greatest in the 1 to 10 cm category. This ranged from 70% at Site 1 to 97% at Site 7. At site 8, 66% of the fish were in the >20 cm category. This was due to an extremely large school of >30 cm *Aethaloperca rogaa* (Serranidae) aggregated on a coral outcrop at 20 m depth.




Figure 22. The pattern of fish species distribution around the outer reef of the atoll in February 2001.






Figure 23. The pattern of fish species distribution around the outer reef of the atoll in November 1999.

There were no significant correlations between the densities of fishes at Sites 1-8, and the east to west positions along the outer reef.<sup>3</sup>

### ***Relationship with Coral Habitat***

There was no significant correlation between the number of species of fish, nor the density of fishes, surveyed at both depths at Sites 1-8, and the amounts of live coral habitat, and live coral and dead coral habitat combined, at each site.

The families Pomacentridae, Chaetodontidae, Labridae, and Serranidae each have several species that are commonly associated with live coral habitat, and habitat structure formed by erect dead corals. These selected fishes were examined for any associations with these measures of habitat structural complexity.

#### ***a) Fish Density***

Only the density of Labrids at 10 m depth ( $R^2 = 0.78$ ,  $F_{.05,1} = 21.33$ ,  $P = 0.004$ ) and at 20 m depth ( $R^2 = 0.54$ ,  $F_{.05,1} = 7.02$ ,  $P = 0.038$ ), and the number of species of Pomacentrids at 20 m depth ( $R^2 = 0.67$ ,  $F_{.05,1} = 12.18$ ,  $P = 0.013$ ), were significantly correlated with the amount of **live coral habitat** at each site. In comparison, both the density of Chaetodontids at 10 m depth ( $R^2 = 0.50$ ,  $F_{.05,1} = 6.01$ ,  $P = 0.050$ ) and 20 m depth ( $R^2 = 0.60$ ,  $F_{.05,1} = 9.08$ ,  $P = 0.024$ ), and the density of Labrids at 10 m depth ( $R^2 = 0.70$ ,  $F_{.05,1} = 14.21$ ,  $P = 0.009$ ) and 20 m depth ( $R^2 = 0.52$ ,  $F_{.05,1} = 6.61$ ,  $P = 0.042$ ), were significantly correlated with the amount of **combined live and dead coral habitat** at each site.

#### ***b) Fish Species***

For the number of species of fish, only the Chaetodontids ( $R^2 = 0.52$ ,  $F_{.05,1} = 6.53$ ,  $P = 0.043$ ) and Pomacentrids ( $R^2 = 0.64$ ,  $F_{.05,1} = 10.73$ ,  $P = 0.017$ ) were significantly correlated with the amount of **combined live and dead coral habitat** at each site.

## **Temperature Data Loggers**

Since the February 2001 deployment of the 10 water temperature data loggers, three have been regularly monitored by Aldabra station staff, and appear to be functioning as anticipated. Although they are developing encrusting growths (Figure 24), it is unlikely that these would interfere with the operation and accuracy of the loggers<sup>4</sup>.

The temperature data from Site 1 (19 February to 20 June 2001) at 10 m depth on the outer reef had a maximum of 29.7 °C (April), a minimum of 21.4 °C (February) and a mean of 27.4 °C for this period (Table 5). There was a significantly decreasing trend in the temperatures at this outer reef site for this period ( $R^2 = 0.59$ ,  $F_{.05,1} = 9818.18$ ,  $P = 0.0$ ; Figure 25).

---

<sup>3</sup> The number of species of fish, and the fish per 100 m<sup>2</sup>, given in Figure 21 correct for errors in some of these numbers for Sites 1, 5, 6, and 7 in Figure 10 in Teleki et al. (1999).

<sup>4</sup> OnSet Computer Corp was consulted about the effect which the encrustations would have on the temperature. Given that the electronics, including the sensor, are enclosed in a waterproof plastic housing, their engineering staff do not believe that external growth would cause changes in the accuracy of the logger's sensor or electronics. The growths will add thermal mass to the housing, which could affect time response. However, the housing is thick, so the expectation is that a fairly substantial amount of growth would be required to affect the response time of the logger.

Figure 24. Encrustation on a temperature logger deployed in this survey.




Figure 25. Temperatures recorded at 30 minute intervals at Aldabra Marine Programme survey Site 1 at 10 m depth, from 18 February to 8 July 2001.




Figure 26. Temperatures recorded at 30 minute intervals at Aldabra Marine Programme temperature monitoring site Passe Dubois at 3 m depth, from 19 February to 19 June 2001.




Figure 27. Temperatures recorded at 30 minute intervals at Aldabra Marine Programme temperature monitoring site North Ile Esprit at 3 m depth, from 11 April 2001 to 10 July 2001.

The temperature data for Passe Dubois (19 February to 20 June 2001) at 3 m depth had a maximum of 32.7 C° (March and April), a minimum of 23.6 C° (June) and a mean of 28.0 C° for this period (Table 5). There was a steadily decreasing trend in maximum and minimum values towards June. Despite the wide fluctuations in temperature on most days, there was a significantly decreasing trend in the temperatures at 3 m in this passage to the lagoon for this period ( $R^2 = 0.33$ ,  $F_{.05,1} = 2883.62$ ,  $P = 0.0$ ; Figure 26).

The temperature data for Île Esprit (11 April to 10 July 200) at 3 m depth had a maximum of 31.2 C° (April), a minimum of 21.8 C° (June) and a mean of 26.8 C° for this period (Table 5). There was a steadily decreasing trend in maximum and minimum values towards July. There was a significantly decreasing trend in the temperatures at 3 m in area of the lagoon for this period ( $R^2 = 0.80$ ,  $F_{.05,1} = 17315.99$ ,  $P = 0.0$ ; Figure 27).<sup>5</sup>

Table 5. Temperatures (°C) recorded, by monthly period, at Aldabra Atoll by the Optic StowAway temperature loggers deployed during the February 2001 AMP expedition (SD = standard deviation, n = number of 30 minute interval recordings). A. Site 1 at 10m depth, 18 February to 8 July; B. Passe Dubois at 3 m depth, 19 February to 20 June; and C. Île Esprit at 3 m depth 11 April – 10 June 2001.

**A. Site 1 at 10m depth**

<b>Date</b>	<b>Mean</b>	<b>Max</b>	<b>Min</b>	<b>SD</b>	<b>n</b>
<b>Feb</b>	28.3	29.3	26.0	0.6	507
<b>March</b>	28.6	29.3	25.7	0.6	1488
<b>April</b>	28.5	29.7	25.7	0.7	1440
<b>May</b>	27.8	28.9	23.5	0.6	1488
<b>June</b>	26.0	27.5	23.5	0.6	1439
<b>July</b>	25.3	25.7	21.4	0.4	364
<b>Overall</b>	<b>27.4</b>	<b>29.7</b>	<b>21.4</b>	<b>1.3</b>	<b>6725</b>

**B. Passe Dubois at 3 m**

<b>Date</b>	<b>Mean</b>	<b>Max</b>	<b>Min</b>	<b>SD</b>	<b>n</b>
<b>Feb</b>	28.7	32.3	26.8	0.9	462
<b>March</b>	29.1	32.7	26.8	1.0	1488
<b>April</b>	28.8	32.7	26.4	1.1	1440
<b>May</b>	27.8	30.8	23.9	1.0	1488
<b>June</b>	26.3	29.7	23.6	0.9	936
<b>Overall</b>	<b>28.0</b>	<b>32.7</b>	<b>23.6</b>	<b>1.4</b>	<b>5814</b>

**C. Île Esprit at 3 m depth**

<b>Date</b>	<b>Mean</b>	<b>Max</b>	<b>Min</b>	<b>SD</b>	<b>n</b>
<b>April</b>	28.9	31.2	25.4	0.9	940
<b>May</b>	27.6	30.8	25.4	0.9	1488
<b>June</b>	25.3	28.2	21.8	0.9	1440
<b>July</b>	24.6	26.1	22.5	0.6	450
<b>April-July</b>	<b>26.8</b>	<b>31.2</b>	<b>21.8</b>	<b>1.8</b>	<b>4318</b>

<sup>5</sup> The temperature logger data was provided by the following Seychelles Islands Foundation personnel at Aldabra Station: Anna Liljevik, Research Officer, Tina Dubel, Ranger, Conrad Savy, Volunteer, and Allen Cedras, Logistics Manager.



Fish of Aldabra – *Acanthurus lineatus* (top), *Epinephelus* sp. (middle), *Platax orbicularis* (bottom)





## Discussion

### Coral Community

The 1999 survey results showed that coral bleaching in 1998 had had a pronounced effect on the reefs around Aldabra Atoll, particularly in shallow water (Teleki *et al.* 1999). Between 1999 and 2001 there has been little change in the outer reefs, with some suggestion of coral recovery at both the depths studied. There is no evidence of high macro algal growth since 1999, with the exception of site 7 and, to a lesser extent, Site 6. These sites had high coral mortality in 1999 providing a new niche for algae, particularly at Site 7 which appears to have provided suitable conditions for abundant coralline algae growth.

By 2001 the dead corals still visible in 1999 had been mostly broken down to rubble and sand, or were covered by other organisms. The shapes of dead massive colonies were still visible in 2001, and some showed evidence of tissue regeneration from surviving areas around the side and base of the colonies. Many are now covered with coralline algae that are known to provide a good substrate for settlement of coral larvae.

The detected natural variation in live coral cover around Aldabra Atoll can be attributed to the prevailing weather and current patterns. The area encompassing Sites 1, 2 and 6 is the most sheltered and has the highest live coral cover. Live coral cover at depth was good around most of the atoll, with the exception of Site 5 that appears to be a rubble zone.

The new sites established in the lagoon have shown that coral cover is high in the vicinity of the drainage channels. Live coral cover at Sites 9 and 10 was estimated to be double that of most outer reef sites, though species diversity appears lower. There is also less evidence of old dead coral colonies in the lagoon, suggesting that the effect of the 1998 bleaching may not have been as great here, a theory supported by observations made by Teleki *et al.* (1998) just after the 1998 bleaching. Lagoon species may be acclimatised to extreme temperature changes, and new tides may bring in deep cooler water through the channels on every tide change reducing the time lagoon corals are exposed to extreme temperatures.

The temperature data recorded by the three loggers during April through June/July 2001 supports the suspected normally wider range in temperatures in the lagoon. During this period the temperature variation at the two lagoon sites was 9.1 and 9.4 °C respectively and at the outer reef site 6.1 °C.

Coral recruitment at Aldabra appears to be good, with most occurring at around 10 m. This would be expected as corals in shallower than 10 m depth have to cope with higher exposure to swell and a mobile substrate making settlement difficult, while corals in deeper water have less available light and reduced available settlement space due to the higher proportion of live coral. Levels of recruitment at Aldabra are in line with those found at other locations, though direct comparison is not always possible due to differing sampling methods (Miller *et al.* 2000). The diversity of coral families recruiting to Aldabra is high and to be expected in view of the high coral diversity on reefs around the atoll.

Size frequencies of recruits measured in 2001 suggests that there may have been several cohorts recruited since the coral bleaching of 1998. It appears that the fast colonisers such as *Acropora* and *Pocillopora* have had the best chance to produce several cohorts due to their high rate of growth and reproduction. The presence of so many recruits and evidence of several cohorts suggests that the atoll may be primarily self-seeding, though it is conceivable that the predominant southeasterly current may be carrying recruits from Assumption, Astove, Cosmoledo and potentially sites at much greater distances afar.

## **Fish Community**

Unlike sessile coral colonies, fish populations are notoriously variable, even when surveyed at short time intervals. The following conclusions are therefore made within the limitations of two surveys having been undertaken some 16 months apart.

Comparing the 1999 and 2001 fish species lists there is generally good agreement between the two surveys even though the total numbers identified vary somewhat: in 1999, 35 families and 211 species and in 2001, 40 families and 205 species. Using a different survey method, Spalding identified 35 families and 287 species in 1998, prior to the bleaching episode.

Given the broadly similar diversity it is therefore of note that a comparison of the actual counts of fishes and species at each site in 1999 and 2001 appear to be very different (Figures 22 and 23). At this stage it is not known whether this difference is attributable to the different seasons in which the surveys were conducted, some structural change in the fish community, or a survey error. It is nevertheless concluded that surveys should, if possible, be conducted at similar times of the year, ideally in November and February.

If one compares the findings of the 1999 and 2001 surveys, the following conclusions emerge:

1. In 1999 the lowest fish density was at Site 5, and the highest at Site 6. In 2001 the highest count was still at Site 6, but the lowest at Site 3. This may be due to an anomaly at Site 5 where a large school of fusiliers crossed the survey area. If this number is excluded, Site 5 yields the lowest count in 2001.
2. In both years most of the fish counted were in the 0 – 10 cm length category. This was due to large numbers of fish from families with numerous small-sized species (Serranidae (fairybasslets), Apogonidae, and Pomacentridae), and because of juvenile



life stages (e.g. Serranidae). Exceptionally at Site 8 most fish were in the >20 cm category.

3. In both years there was no correlation between the **densities** of fishes counted and the relative positions of the survey sites east to west along the atoll.
4. In 1999 there was also no correlation between the **number of species** counted and the east/west position of the survey sites, but in 2001 a significant correlation did appear, both at the 10 m ( $R^2 = 0.41$ ) and 20 m depths ( $R^2 = 0.47$ ).
5. On coral reefs, fish diversity and habitat complexity are often linked (Ebeling and Hixon 1991; Sebens 1991; Turner *et al.* 1999; Williams 1991). In both 1999 and 2001 a distinction was made between benthic habitat whose structural integrity was still evident (live or dead coral), and habitat without such structural integrity (sand, rock and rubble). In 1999 no correlation was found between the **density** of fish, nor the **number of species**, and either the live or dead coral cover. In 2001 the analysis was slightly different, but a similar conclusion was drawn.
6. However, in 1999 an analysis of fish families that have species commonly associated with structurally complex habitat concluded that there were positive correlations amongst the Chaetodontidae, Labridae and Serranidae and live or dead coral. Although this association was not found amongst the Serranidae in 2001, it remained within the Chaetodontidae and Labridae, and was evident amongst the Pomacentridae at 20 m depth.

Although it is too early to establish any significant trends in Aldabra's fish populations, the nature of any long term study means that these will emerge in time. In particular, close attention should continue to be paid to the relationship between coral reef related species and reef structure. For example, the numbers of specific corallivorous Chaetodontids may prove to be an effective indicator of reef health (Crosby and Reese 1996).

## Temperature

The temperature data for the marine environment at Aldabra being recorded by the loggers deployed during the February 2001 AMP expedition has a great potential to yield a valuable long-term data base for the atoll. This is based on the assessment of the temperature data provided by the Aldabra Station monitoring of three of the loggers, and by the seemingly flawless performance of these three loggers. The temperatures recorded at Site 1 at 10 m depth (Figure 25) show February through June 2001 monthly averages (Table 5) that track favourably with the monthly mean temperatures for these months for the period 1961-1996 (Figure 4). The potential for the encrusting growths developing on the loggers to affect the temperature readings, while suspected not to be a problem, will be tested on the next AMP expedition to Aldabra.

## The Aldabra Marine Programme: Short Term and Long Term

The Aldabra Marine Programme has now established eleven permanent monitoring sites around Aldabra, seven of which have now been surveyed twice. These sites provide baseline data and will generate increasingly more important data as they are resurveyed on a yearly

basis. Furthermore, as the information is gathered and analysed more questions about the atoll are generated, and therefore the scope to amplify the study.

After completing the 1999 survey, the AMP set out a list of five goals for the future. All of these goals have been achieved with the exception of the initiation of a study of mangrove ecosystem dynamics. AMP has therefore increased its research base in 2001 by:

- Continuing the fish and coral monitoring of permanent sites, and establishing one new long term site on the highly exposed southern shoreline;
- Establishing permanent monitoring sites in the lagoon and Passe Houareau channel;
- Initiating a coral recruitment study in the lagoon and outer reef slopes, and tagging corals to follow their survival and growth;
- Deploying temperature data loggers;

There was also the opportunity to train Aldabra staff in survey techniques for coral recruitment and echinoderm population assessment, and to discuss suitable monitoring programmes for the station staff. AMP remains committed to training Seychellois rangers in marine survey techniques and hopes to continue doing so in the future.

A long term marine science programme for Aldabra has now been established by AMP. Future surveys will continue to monitor the permanent sites and it is hoped that another site will be set up on the southern coastline along with further lagoon sites. The team also aims to improve the scope of the physical marine monitoring and expand the lagoon work, hopefully by appointing a PhD student.

## Acknowledgements

We would like to thank the following individuals and organisations for their invaluable support and advice.

Phase II of the Aldabra Marine Programme trip would not have been possible without the generous support from the following organisations and individuals: Coral Reef Degradation in the Indian Ocean (CORDIO) Programme (Olof Lindén), The Global Environment Facility, *The Shoals of Capricorn Programme*, The Royal Geographical Society, The Royal Society, and British Airways Assisting Conservation.

Guillermo Cryns continues to provide us with unwavering support and steadfast dedication to the long term success of the Aldabra Marine Programme.

The following individuals gave AMP generous financial support: Maya Moltzer, Mallorca, Spain; Frank Smith, Pinney Allen, Robyn Ice, T. McCarthy and S. Train (Alston & Bird LLP, Atlanta, USA)

The following people and groups are thanked for their advice, support in the field and assistance to the Aldabra Marine Programme:

Etienne Quilindo who was part of the AMP Phase II research team provided dedicated support and field assistance which allowed AMP to achieve all of its objectives.

Rolph Payet, Seychelles Ministry of Environment and Transport; David Rowat and the Indian Ocean Explorer; Patrick Lablache, Ministry of Land Use and Habitat; Seychelles Island Foundation: Executive Director Lindsay Chong-Seng and Executive Officer Angela Valente-Libanotis; Aldabra Station: Guy Esparon - Manager, Anna Liljevik - Research Officer, Tina Dubel - Ranger, Conrad Savy - Volunteer, Allen Cedras - Logistics Manager, Ross Wanless - Researcher and Brian Betsy – Ranger; David Boullet, Edwin Grancourt and the Seychelles Fishing Authority.

Maurice Loustau-Lalanne SIF Chairman and Principal Secretary, Seychelles Ministry of Environment and Transport; Patrick Lablache, Seychelles Ministry of Land Use and Habitat; Sue Wells, IUCN East Africa Region Office; Stephen Blackmore, Royal Botanic Gardens, Edinburgh UK; and Claude Pavard, Photographer.

Other participants of the Aldabra Scientific Workshop (December 2000) whose advice, suggestions and support for AMP have been and continue to be invaluable: Katy Beaver, Roger Bour, David Bourn, Joseph Francois, Justin Gerlach, Jeanne Mortimer, Robert Prys-Jones, Selby Remie and Richard White

Jeanne Mortimer continues to be a wealth of support and assistance both scientifically and with the logistics of conducting marine research on Aldabra.

The *Shoals of Capricorn* Seychelles based team (Martin Callow, Jan Robinson and Caroline Lawton) provided us with invaluable support and hospitality on Ste. Anne. John Collie and the staff of the Marine Parks Authority were very accommodating in transporting us to and from Victoria, helping us shift equipment, and allowing us to have access to the facilities of Ste. Anne.

Tom Spencer, *Shoals of Capricorn* Science Plan Co-ordinator, provided us with valuable advice on both the logistics and the science plan of the research programme. His comments on the final report proved insightful and invaluable. Juliet Burnett and Jessica Kavanagh of the *Shoals of Capricorn*, RGS London Office were outstanding in coordinating the overall logistics in the UK and providing valuable contacts in the Seychelles.

David Stoddart, Peter Sale, Callum Roberts and Barbara Brown were generous with their time in reviewing funding proposals and providing comments on the AMP science plan.

Clare Bradshaw, Mark Spalding, Tom Spencer and Kristian Teleki made important contributions as members of the original Southern Seychelles Atoll Research Programme (SSARP) who first visited Aldabra in 1998 at the peak of the coral bleaching event.

The Cambridge Coastal Research Unit (CCRU) has provided immense support financially and logistically, and by allowing their personnel to participate in the expedition. Fabrice Meziani, Eureka Design Ltd., kindly helped with some of the artwork in this report.

Camera Pix, Nairobi generously donated an aerial photograph of Aldabra which was used on both the cover of this report and on the AMP brochure. Alan Downing also kindly donated his time to edit the text of the AMP brochure.

The enthusiasm, encouragement and support from Professor David Stoddart, who is the reason why any scientific research can be conducted on Aldabra, were deeply appreciated.

# References

- Arnoult, J., M.L. Bauchot-Boutin and R. Roux-Estève. 1958. Les poissons de l'Île Aldabra. *Annales de l'Institut Océanographique* 34:47-90.
- Attenborough, D. 1995. Foreword. In M. Amin, D. Willetts, and A. Skerrett, (eds) *Aldabra World Heritage Site*. Camerapix Publishers International, Nairobi, Kenya. p 7.
- Barnes, J., D.J. Bellamy, D.J. Jones, B. Whitton, E. Drew, L. Kenyon, J.N. Lythgoe and B.R. Rosen. 1971. Morphology and ecology of the reef front of Aldabra. *Symposium Zoological Society London* 28: 87-114.
- Beaver, K. and R. Gerlach. 1997. *Aldabra Management Plan: a management plan for Aldabra Atoll, Seychelles Natural World Heritage Site 1998-2005*. Seychelles Island Foundation and the GEF/World Bank. p 61.
- Birkeland, C. 1997. Symbiosis, fisheries and economic development on coral reefs. *Trends in Ecology and Evolution* 12:364-366.
- Bohnsack, J.A. 1995. Maintenance and recovery of reef fish productivity. In N.V.C. Polunin and C.M. Roberts (eds), *Management of Reef Fisheries*. Chapman & Hall, London. p 283-313.
- Bohnsack, J.A. and J.S. Ault. 1996. Management strategies to conserve marine biodiversity. *Oceanography* 9(1):73-82.
- Brown, B.E. 1997. Coral bleaching: causes and consequences. *Coral Reefs* 16:S129-S138.
- Brown, B.E., R.P. Dunne, and H. Chansang. 1996. Coral bleaching relative to elevated seawater temperature in the Andaman Sea (Indian Ocean) over the last 50 years. *Coral Reefs* 15:151-152.
- Bryant, D., L. Burke, J. McManus and M. Spalding. 1998. *Reefs at risk: a map-based indicator of threats to the world's coral reefs*. World Resources Institute, Washington, D.C. p. 56.
- Chadwick, D.H. 1999. Coral in peril. *National Geographic* 195(1):30-37.
- Crosby, M.P. and E.S. Reese. 1996. A manual for monitoring coral reefs with indicator species: butterflyfishes as indicators of change on Indo Pacific reefs. Office of Ocean and Coastal Resource Management, National Oceanic and Atmospheric Administration, Silver Spring, MD. p. 45
- Davies, J.M., R.P. Dunne and B.E. Brown. 1997. Coral bleaching and elevated seawater temperature in Milne Bay Province, Papua New Guinea, 1996. *Marine and Freshwater Research* 48: 513-516.
- Doubilet, D. 1999. Coral eden. *National Geographic* 195(1):2-29.
- Drew, E.A. 1977. A photographic survey down the seaward reef-front of Aldabra Atoll. *Atoll Research Bulletin* 193.
- Ebeling, A.W. and M.A. Hixon. 1991. Tropical and temperate reef fishes: comparisons of community structures. In P.F. Sale (ed) *The Ecology of Coral Reef Fishes*. Academic Press, San Diego, California. p 509-563.
- Engelhardt, U. 2001. Ecological characteristics of scleractinian coral communities around the inner granitic islands of the Seychelles 2 1/2 years after the 1998 mass coral bleaching event. Reefcare International Pty Ltd, Townsville, Australia. p. 86.
- English, S., C. Wilkinson and V. Baker. 1997. *The survey manual for tropical marine resources*. 2<sup>nd</sup> ed. Australian Institute of Marine Science. P. 390.
- Ginsburg, R.N., P. Kramer, J. Lang, P. Sale and R. Steneck. 1998. Revised rapid assessment protocol (RAP). Atlantic and Gulf Reef Assessment (AGRA). <http://coral.amol.noaa.gov/agra/rap-revised.html>.
- Glynn, P.W. and W.H. Deweerdt. 1991. Elimination of two reef building hydrocorals following the 1982-83 El Niño warming event. *Science* 253:69-71.
- Hodgson, G. 1999. A global assessment of human effects on coral reefs. *Marine Pollution Bulletin* 38(5):345-355.
- Jatzow, R. and H. Lenz. 1899. *Fische von Ost-Afrika, Madagaskar und Aldabra*. *Abhandlungen der Senckenbergischen naturforschenden Gesellschaft* 21:497-531.

- Johannes, R.E. 1981. Words of the Lagoon. University of California Press, Berkeley, p. 320.
- McPhaden, M.J. 1999. The child prodigy of 1997 - 98. *Nature* 398: 559-562.
- Miller, M.W., Weil, E., and Szmant, A.M. 2000. Coral recruitment and juvenile mortality as structuring factors for reef benthic communities in Biscayne National Park, USA. *Coral Reefs* 19: 115-123.
- Mortimer, J.A. 1988. Green turtle nesting at Aldabra Atoll - population estimates and trends. *Bulletin of the Biological Society of Washington* 8:116-128.
- Mortimer, J.A. 1997. Turtle monitoring at Aldabra (1997 Version). Global Environmental Facility (EMPS Project J1: Turtle and Tortoise Conservation) and the Government of the Seychelles. P. 47.
- Osborne, K. and W.G. Oxley. 1997. Sampling benthic communities using video transects. In S. English, C. Wilkinson and V. Baker (eds) *Survey Manual for Tropical Marine Resources*, 2<sup>nd</sup> Edition. Australian Institute of Marine Science, Cape Ferguson. p. 363-376.
- Polunin, N.V.C. 1984. Marine fishes of the Seychelles. In D.R. Stoddart (ed) *Biogeography and ecology of the Seychelles Islands*. W. Junk Publishers, Netherlands. p 171-191.
- Potts, G.W. 1973. The ethology of *Labroides dimidiatus* (Cuv. & Val.) (Labridae, Pisces) on Aldabra. *Animal Behaviour* 21:250-291.
- Regan, C.T. 1912. New fishes from Aldabra and Assumption collected by Mr. J.C.F. Fryer. *Percy Sladen Expedition Reports* 4. *Transactions of the Linnean Society, London Ser. 2, Zoology* 15:301-302.
- Roberts, C.M., J.P. Hawkins, D.E. McAllister and F.W. Schueler. In press. Hotspots, endemism and the conservation of coral reef fish biodiversity. *Coral Reefs*.
- Roberts, C.M., J.P. Hawkins, N. Chapman, V. Clarke, A.V. Morris, R. Miller and A. Richards. 1999. The threatened status of marine species. Center for Marine Conservation/IUCN, Washington, D.C.
- Robertson, D.R., N.C.V. Polunin and K. Leighton. 1979. The behavioral ecology of three Indian Ocean surgeonfishes (*Acanthurus lineatus*, *A. leucosternon* and *Zebrasoma scopas*): their feeding strategies, and social and mating systems. *Environmental Biology of Fishes* 4(2):125-170.
- Russ, G.R. 1991. Coral reef fisheries: effects and yields. In P.F. Sale (ed.) *The Ecology of Coral Reef Fishes*. Academic Press, San Diego, California. Chapter 20, p. 601-635.
- Sale, P.F. 1991. Introduction. In P.F. Sale (ed) *The Ecology of Coral Reef Fishes*. Academic Press, San Diego, California, p. 3-15.
- Sebens, K.P. 1991. Habitat structure and community dynamics in marine benthic systems. In S.S. Bell, E.D. McCoy and H.R. Mushinsky (eds) *Habitat Structure – The Physical Arrangement of Objects in Space*. Chapman and Hall, London. p. 211-234.
- Shapiro, D.Y. 1977. Social organization and sex reversal of the coral reef fish *Anthias squamipinnis* (Peters). University of Cambridge Ph.D. Thesis. p. 386.
- Skerrett, J. and L. Mole. 1995. Somewhere in an empty ocean. In M. Amin, D. Willetts, and A. Skerrett, (eds.) *Aldabra World Heritage Site*. Camerapix Publishers International, Nairobi, Kenya. p. 23-48.
- Slingo, J. (1998). The 1997/1998 El Niño. *Weather* 53: 274-281.
- Smith, J.L.B. and M.M. Smith. 1969. *Fishes of the Seychelles* (2<sup>nd</sup> edition). Rhodes University Institute of Ichthyology, Grahamstown. p. 215.
- Spencer T., K.A. Teleki, C. Bradshaw and M.D. Spalding. 2000. Coral bleaching in the Southern Seychelles during the 1997-1998 Indian Ocean warming event. *Marine Pollution Bulletin* 40(7): 569-586.
- Steneck, R.S. 1998. Human influences on coastal ecosystems: does overfishing create trophic cascades? *Theoretical Ecology* 13(11):429-430.
- Stevens, J.D. 1984. Life-history and ecology of sharks at Aldabra Atoll. *Proceedings Royal Society of London B* 222:79-106.

- Stoddart, D.J. 1984. Coral reefs of the Seychelles and adjacent islands. In D.R. Stoddart, (ed) Biogeography and ecology of the Seychelles Islands. W. Junk Publishers, Netherlands. pp 31-68
- Stoddart, D.J. and L.U. Mole. 1977. Climate of Aldabra Atoll. Atoll Research Bulletin 202.
- Stoddart, D.R. 1971. Scientific studies of Aldabra and neighbouring islands. Philosophical Transactions of the Royal Society of London B. 260:5-29.
- Stoddart, D.R. 1983. Spatial and temporal variability of rainfall on Aldabra Atoll. Atoll Research Bulletin 273.
- Stoddart, D.R. 1997. Bibliography of Aldabra Atoll, 3<sup>rd</sup> Edition. Seychelles Island Foundation, London and Mahe. p. 89.
- Teleki, K.A., N. Downing, B. Stobart and R. Buckley. 2000. Aldabra: Marine monitoring and management into the 21<sup>st</sup> century. Cambridge Coastal Research Unit, Department of Geography, University of Cambridge. p. 23.
- Teleki, K.A., N. Downing, B. Stobart and R. Buckley. 1999. Aldabra Marine Programme. Cambridge Coastal Research Unit, Department of Geography, University of Cambridge. p. 31.
- Teleki, K.A., T. Spencer, C. Bradshaw and M.D. Spalding. 1998. Coral Bleaching in the Western Indian Ocean – a sign of the times? International Society for Reef Studies – European Meeting, 1-4 September, Ecole Pratique des Hautes Etudes, Universite de Perpignan.
- Turner, J.S., S.F. Thrush, J.E. Hewitt, V.J. Cummings and G. Funnell. 1999. Fishing impacts and the degradation or loss of habitat structure. Fisheries Management and Ecology 6:401-420.
- Van Moorsel, G.N.M. (1988) Early maximum growth of stony corals (Scleractinia) after settlement on artificial substrata on a Caribbean reef. Marine Ecology Progress Series 50: 1008-1009.
- Viles, H., T. Spencer, K. Teleki and C. Cox. 2000. 16 years of microfloral recolonisation data from limestone surfaces, Aldabra Atoll, Indian Ocean: Implications for biological weathering. Earth Surface Processes and Landforms. 25(12): 1355-1370.
- Wallace C.C. 1985. Reproduction, recruitment and fragmentation in nine sympatric species of the coral genus *Acropora*. Marine Biology 88: 217-233.
- Wallace, C.C., A. Watt and G.D. Bull. 1986. Recruitment of juvenile corals onto coral tables preyed upon by *Acanthaster planci*. Marine Ecology Progress Series 32: 299-306.
- Wilkinson, C.R. 1998. The 1997-1998 mass bleaching event around the world. In C.R. Wilkinson (ed.) Status of Coral Reefs of the World: 1998. Australian Institute of Marine Science, Cape Ferguson. p. 15-38.
- Wilkinson, C.R. 2000. Status of coral reefs of the world:2000. Australian Institute of Marine Science, Cape Ferguson. p. 363.
- Wilkinson, C.R. and R.W. Buddemeier. 1994. Global climate change and coral reefs: implications for people and reefs. Report of the UNEP-IOC-ASPEI-IUCN Global Task Team on the Implications of Climate Change on Coral Reefs. IUCN, Gland Switzerland. p. 124.
- Williams, D.McB. 1991. Patterns and processes in the distribution of coral reef fishes. In P.F. Sale (ed) The Ecology of Coral Reef Fishes. Academic Press, San Diego, California. p. 437-474.
- Zar, J.H. 1984. Biostatistical analysis, 2<sup>nd</sup> edition. Prentice-Hall, Inc., New Jersey.